

of it, but shall hope to greet it again another year. Gray's Manual gives the range of this species as from Illinois and Missouri to North Carolina and Texas. It has, however, been known for some time near Providence, Rhode Island (see W. W. Bailey, RHODORA, iv. 198), and I learn from the Gray Herbarium that Mr. J. C. Parlin has found it spreading from wool-waste at North Berwick, Maine.

It seems probable that it may have been introduced at the Connecticut station in grass seed.—FRANCES M. GRAVES, New London, Connecticut.

---

FURTHER STATIONS FOR BOTRYCHIUM MATRICARIAEFOLIUM IN CONNECTICUT.—Bishop's list gives only two stations for *Botrychium matricariaefolium*, one in Sherman (Fairfield County), the other in Franklin (New London County). In RHODORA, 3: 36 (1901) Mr. A. W. Driggs notes the finding of *B. matricariaefolium* in West Hartford (Hartford County). In June, 1901, the writer collected several specimens at Mansfield (Tolland County). During the summer of 1902 at Cornwall Bridge (Litchfield County) many specimens were found; and at Kent Falls, North Kent, in the same county one plant was collected. Last summer (1903) this species was found to be abundant on the mountain slopes in Salisbury (Litchfield County). Specimens from these stations are preserved in the writer's herbarium.

As this plant has been found in five of the eight counties and in widely separated parts of the state, it is probable that it has been overlooked and will be found to be generally distributed throughout.—A. VINCENT OSMUN, Amherst, Massachusetts.

Vol. 6. no. 63, including pages 45 to 64, was issued 7 March, 1904.